Unleashing the Power of Alkali-Metal Mediation for Arene Functionalisation and Catalysis



Eva Hevia

Ischia Advanced School of Organic Chemistry


19th September 2026

UNIVERSITÄT BERN

Organolithium Reagents, the Workhorses of Synthesis



W. Schlenk,
Ber. Dtsch. Chem. Ges.
1917, *50*, 262




Efavirenz (Antiretroviral)

(€3.11 billion per year)

- ✓ vast uses in synthesis
- ✓ commercially available
- ✓ high reactivity
- ✓ soluble in hydrocarbon solvents






$[^{n}BuLi]_6$

D. Stalke,
Angew. Chem, Int. Ed.
1993, *32*, 580


Heavier Alkali-Metal Reagents (NaR, KR)




J. A. Wanklyn,
Proc. R. Soc.
1858, *9*, 341



Abundance: 0.002 %
Electronegativity: 0.98
Size: 145 pm



Abundance: 2.6 %
Electronegativity: 0.93
Size: 180 pm



Abundance: 2.4 %
Electronegativity: 0.82
Size: 235 pm


Main Limitations of NaR, KR

- **Lack of solubility**
- **Too reactive**
- **Lack of thermal stability**
- **non-commercially available**

Renaissance of MR (M = Na, K) in Organic Chemistry


UNIVERSITÄT BERN

Na dispersion



S. Asako, K. Takai, L. Ilies, *Nat.Cat.* **2019**, *2*, 297;
Adv. Synth. Cat. **2019**, *361*, 3120; *Commun. Chem.* **2021**, *4*, 76

Na-packed bed reactor




P. Knochel, *Angew. Chem. Int. Ed.* **2021**, *60*, 14296;
Angew. Chem. Int. Ed. **2022,** *61,* e202203807

NaAr intermediates are prepared *in situ*!

Potassium Mediated Zincation Reactions


Metalation of fluoroarenes



Lochmann-Schlosser "LICKOR" Superbase


M. Schlosser, *Tetrahedron Lett.* **1996**, *36*, 6551

$Zn(TMP)_2$






0.5M solution
in toluene




no reaction



>99% (76%)



Neil Judge




thermally stable intermediate
Zn-C vs K-F bonding
higer order zincate (2:1 K:Zn)

direct Zn-H exchange
two fold activation of
$Zn(TMP)_2$!


Angew. Chem. Int. Ed. **2023**, *63*, e202303099

Potassium Mediated Zincation of Fluoroarenes




Yields determined by ^{1}H NMR spectroscopy, isolated yields shown in brackets


polyfluorinated aromatics



10 min, >95%
(76%)




10 min, >95%
(70%)




10 min, >95%
(64%)

non-activated fluoroarenes




24 h, >95%
(65%)




24 h, 82%
(75%)

p-substituted halofluoroarenes




1 h, >95%
(72%)




1 h, >95%
(73%)


fluorotoluenes



24 h, >95%
(85%)




24 h, >95%
(81%)




24 h, >95%
(70%)

fluoroanisoles




24 h, >95%
(57%)




24 h, 90%
(67%)


miscellaneous fluoroarenes



24 h, 93%
(76%)




3 h, >95%
(67%)



10 min, >95%
(78%)

Angew. Chem. Int. Ed. **2023**, *63*, e202303099

Potassium Mediated Zincation of Fluoroarenes




Angew. Chem. Int. Ed. **2023**, *63,* e202303099

Potassium Mediated Zincation of Non-activated Substrates

UNIVERSITÄT BERN


$Zn(TMP)_2$

commercially available
0.5M solution in toluene







30 min, 95% 30 min, 89%



Zn-σC vs K-πC bonding






drastic alkali-metal effect:
✗ Li, Na ✓ K

Benzylic aroylation of toluene

zincate modification:
enhanced basicity and Nu power


mechanistic studies




active base:
potassium zincate

Chem. Sci. **2024**, *15*, 14757


For a Tutorial Review on this topic:
Chem. Soc. Rev. **2026**, *55*, 8382





Assessing Directed-*ortho*-Sodiation of Anisole





Traditional non-nucleophilic strong bases:



TMEDA  34%


PMDETA  36%






[NaTMP]$_3$

M. F. Lappert, R. E. Mulvey
J. Organomet. Chem., **1999**, *587*, 88





[NaTMP·TMEDA]$_2$


C. T. O'Hara,
Chem. Eur. J., **2008**, *14*, 8025

dimeric structure retained in solution




Andreu Tortajada


Assessing Directed-*ortho*-Sodiation of Anisole




- mixed-aggregate intermediate
- structure retained C_6D_{12} solution
- **43%** metalation




^{1}H-DOSY NMR, C_6D_{12}, 300.1 MHz, 298 K




Sodium Mediated Borylation Approaches





BX_3 — *aryl trapping equilibrium shift*

$B(OiPr)_3$ 83% yield



$B(OMe)_3$ 10% yield

$B(OMe)_3$ reacts with [NaTMP·PMDETA]$_2$


base degradation


steric incompatibility of $B(OiPr)_3$ with [NaTMP·PMDETA]$_2$

LiTMP, 13% KTMP, 5%


Angew. Chem. Int. Ed. **2022**, *61*, e202204262
Chem. Sci. **2023**, *14*, 6538

Applications in Suzuki-Miyaura Couplings







79%




74%




76%




58%




64%




63%
(-78 ºC)




86%
(-78 ºC)




67%
(-78 ºC)



76%
(-78 ºC)




67%
(-78 ºC)



89%



71%




83%

Limitations

- ✗ Cl, Br, CO_2R substituents
- ✗ low T (-78ºC) for ArF
- ✗ unsuccessful for diborylations
- ✗ no compatible with pyridines

Angew. Chem. Int. Ed. **2022**, *61*, e202204262; *Chem. Sci.* **2023**, *14*, 6538


Using an Amino(imino)borane as a Trapping Agent



amino(imino)borane


H. Nöth, *Eur. J. Inorg. Chem.* **2004**, 3612

sp-hybridised B: highly electrophilic
bulky substituents: precude oligomerisation



Clarence Tan

Using an Amino(imino)borane as a Trapping Agent





amino(imino)borane

Na boryl(amide)


95% yield

H. Nöth, *Eur. J. Inorg. Chem.* **2004**, 3612






Clarence Tan




monomer/dimer equilibrium in C_6D_6 solution

(^{1}H DOSY NMR)





Upgrading Deprotonative Borylation to Catalytic Regimes




Sodium-Catalysed Deprotonative Borylation!

LiTMP, 6%
NaHMDS, 34%









J. Am. Chem. Soc.,**2025**, *147*, 42076

Upgrading Deprotonative Borylation to Catalytic Regimes






Key advantages

- ✓ Compatible with Cl, Br substituents
- ✓ Controlled di-borylation
- ✓ Room temperature borylation ArF
- ✓ Remarkable stoichiometric control


J. Am. Chem. Soc.,**2025**, *147*, 42076

Upgrading Deprotonative Borylation to Catalytic Regimes




Iterative Borylations


n =2



88%




94%




48%

n = 1




24%

n =2




95%

n =3




71%




excellent stoichiometric control


J. Am. Chem. Soc.,**2025**, *147*, 42076

Upgrading Deprotonative Borylation to Catalytic Regimes







90%




68%




53%




78%




48%



54%




95%




78%

Catalytic deprotonative borylation of pyridines



J. Am. Chem. Soc.,**2025**, *147*, 42076


Mechanistic Understanding



Kinetic investigations
(initial rates and DFT simulation)
First order in [benzothiophene]
First order in [NaTMP]
Zeroth order in [Boron Trap]

ωB97XD:
6-31+G(d,p)/6-311+G(d,p)

Kinetic Isotopic Effect
KIE = 8.7




TMP(H) is not involved in catalytic cycle!




DFT calculations
Max Garcia-Melchor
(CIC energiGUNE, Spain)

J. Am. Chem. Soc.,**2025**, *147*, 42076


Mechanistic Understanding



Stoichiometric Polyborylation of Non-activated Arenes




40%, 24h, 70°C
sluggish towards
non-activated arenes




1,4-diborylation

2,6-diborylation




44%
(2 equiv NaTMP)

1,1',3,3'-tetraborylation




90%
(5 equiv NaTMP)

2,5,8,11-tetraborylation




41%
(4 equiv NaTMP)

Sodium-mediated Silylation of Arenes and Heteroarenes



Deviation from the standard conditions	Yield [%]
None	73
LiTMP instead of NaTMP	18
$Na(CH_2SiMe_3)$ instead of NaTMP	0
Me_3SiCl instead of Et_3SiCl	0
10 mol% of PMDETA	**84**

Accessing gem-disilylation of toluene




suggested by DOSY-NMR studies



David Sanchez-Roa


Angew. Chem. Int. Ed. **2026**, *65, e1774364*

Sodium-mediated Silylation of Arenes and Heteroarenes




Benzylic C(sp^3)–H silylation


Gram-scale preparation (6 mmol)




1a: 95% (1.17 g)




1b: 79%




1c: 86%




1d: 67%




1e: 84%




1f: 77%




1g: 76%



1h: 71%



1i: 65%




1j: 68%



David Sanchez-Roa


Angew. Chem. Int. Ed. **2026**, *65, e1774364*

Sodium-mediated Silylation of Arenes and Heteroarenes




X = N, O, S **1** **2** **3**

Arenes: C(sp^2)–H silylation



2a: 75% **2b**: 71% **2c**: 71% **2d**: 91% **2e**: 71% **2f**: 72% **2g**: 62%

Heteroarenes: C(sp^2)–H silylation



3a: 92% **3b**: 91% **3c**: 87% **3d**: 83% **3e**: 40%




3f: 76% **3g**: 74% **3h**: 82% **3i**: 74%

Mechanistic Studies: DFT Silylation Pathway

UNIVERSITÄT BERN

Computational studies in collaboration with Prof. Max García Melchor



Angew. Chem. Int. Ed. **2026**, *65*, *e1774364*


Mechanistic Studies: DFT Silylation Pathway

NMR monitoring experiments:

Me_3SiCl: *rapid and complete decomposition*


Et_3SiCl: *slow (partial) decomposition*

iPr_3SiCl: *unreactive towards NaTMP*




Angew. Chem. Int. Ed. **2026**, *65, e1774364*

Decomposition studies (NMR + MS)




Stoichiometric Control in Multisilylation Reactions




X = N, O, S

4


Benzylic C(sp^3)–H multisilylation




4b: 97%




4c: 81%



4d: 96%




4e: 81%




4f: 93%


(Hetero)arenes: C(sp^2)–H multisilylation




4g: 95%




4h: 93%




4i: 88%




4j: 98%



4k: 83%




4l: 95%




4m: 59%


Orthogonal multisilylation




4n: 97%[b]



4o: 97%




4p: 74%




4q: 83%

Angew. Chem. Int. Ed. **2026**, *65, e1774364*


Metalation of Arenes with NaTMP: an Equilibrium



Metalation of Arenes with NaTMP: an Equilibrium




Benzene C-H Deprotonation for Isotopic Incorporation







Andreu Tortajada




Benzene C-H Deprotonation for Isotopic Incorporation







Benzene C-H Deprotonation for Isotopic Incorporation



Assessing Base and Donor Effects on the Deuteration of Anisole





Base	Donor	Time	Degree of deuteration /%			
			D_o	D_m	D_p	OCD_3
NaTMP	PMDETA	4h	>95	>95	>95	>95




NaTMP TMEDA PMDETA

Substrate Scope of NaTMP/PMDETA Catalysed HIE







1a-d_8




1b-d_{10}




1c-d_8




1d-d_5




1e-$d_{11/12}$




1f-d_5




1g-d_6




1h-d_5



1i-d_3




1j-d_9




1k-d_{18}


J. Am. Chem. Soc. **2022**, *144*, 20237; *Cat. Sci. Tech.* **2023**,*13*, 4919

Substrate Scope of NaTMP/PMDETA Catalysed HIE







1l-d_{10}




1m-d_{10}




1n-$d_{9/10}$




1o-d_{12}




1p-d_{12}




1q-d_3/d_6




1r-d_6/d_{10}




1s-d_8



1t-d_{10}




1u-d_4/d_7



1v-d_8/d_{10}


J. Am. Chem. Soc. **2022**, *144*, 20237; *Cat. Sci. Tech.* **2023**,*13*, 4919


Investigating the Role of NaTMP/TMP(D)



no apparent deuteration without TMP


NaTMP / TMP(D) as a shuttle for isotopic exchange!






J. Am. Chem. Soc. **2022**, *144*, 20237; *Cat. Sci. Tech.* **2023**,*13*, 4919

Redesign of the Deuteration Platform








Andreu Tortajada




Melina Tschopp


Redesign of the Deuteration Platform




Heterocycles




NHCs



Fluoroarenes




Amides



Angew. Chem. Int. Ed. **2025**, *64*, e202421736

Brucine (alkaloid)


Redesign of the Deuteration Platform




NaHMDS : **37% D**
$NaCH_2SiMe_3$: **68% D**

HMDS(H) is **not needed** for the HIE
$NaCH_2SiMe_3$ is a **more poweful** catalyst!

Proposed mechanism







catalytically competent

Angew. Chem. Int. Ed. **2025**, *64*, e202421736

Redesign of the Deuteration Platform




16 examples, regioselective deuteration




Angew. Chem. Int. Ed. **2025**, *64*, e202421736

Two Complementary Na-catalysed HIE Methods




✔ *Simple Reagents* ✔ *Mild Conditions* ✔ *Mechanistic Understanding* ✔ *Selective Deuteration*




- ✔ *Efficient Perdeuteration*
- ✔ *Mild Conditions*
- ✔ *Mechanistic Understanding*
- ✔ *Non-activated Substrates*


Deuteration of Cyclohexane: A Gateway for Alkene Isomerisation

deuteration and isomerisation





86%, 90 min; 94:6 *E:Z* selectivity



TMEDA

PMDETA

LiTMP/PMDETA, 7%


NaTMP/TMEDA, 61% (81:19 *E:Z*)


alkali-metal and Lewis donor effects!


Angew. Chem. Int. Ed. **2024**, *63*, e202407262

Mechanistic investigations

Isolation of π-allyl intermediate







TMP(H) accelerating effect!

88%, **3 days**

86:14 *E:Z* selectivity

---


vs NaTMP (PMDETA) cat:

86%, **90 min**

94:6 *E:Z* selectivity

Angew. Chem. Int. Ed. **2024**, *63*, e202407262


Sodium Catalysed Alkene Isomerisation




ωB97XD:6-31+G(d,p)/6-311+G(d,p)


Angew. Chem. Int. Ed. **2024**, *63*, e202407262

Sodium Catalysed Alkene Isomerisation






Limitation:




Mixture of isomers
100% conv.

Angew. Chem. Int. Ed. **2024**, *63*, e202407262

Highly Reactive Hydrocarbon Soluble Alkylsodium Reagents



$NaCH_2SiMe_3$




- thermally stable (lack of of β-Hs)
- $SiMe_3$ can enhance solubility
- highly aggregated structure $[\{Na(CH_2SiMe_3)\}_4]_n$


T. Shun et al, *Chem Commun.* **1993**, 1386

hexane




PMDETA






hexane, RT


$[\{Na(CH_2SiMe_3)\}_4]_n$




^{1}H, ^{13}C and ^{1}H DOSY NMR

Me_6-TREN hexane, RT








Angew. Chem. Int. Ed. **2023**, *62*, e202218498,




David Anderson


Donor-free hydrocarbon soluble alkyl sodium




$[NaCH_2tBu]_4$


- discrete tetramer
- stabilised by dispersion
- soluble in hexane
- thermally stable





MR	Yield (%)
$NaCH_2SiMe_3$	21
$NaCH_2tBu$	**83**
$LiCH_2tBu$	12
$LiCH_2tBu$ + NaOtBu	49







***meta*-regioselectivity**

C-H metalation of **non-activated** substrates

in preparation



Making Organosodium Chemistry Even More Sustainable



Halogen–sodium exchange of aryl bromides


Reductive sodiation of aryl chlorides




Laurean Ilies



Sobi Asako





Asako et al, *Nat. Cat.* **2019**, *2*, 297



FeX_n	Cosolvent	Yield (%)
$FeCl_2$	C_6H_6	1
$FeBr_2$	C_6H_6	5
$Fe(acac)_2$	C_6H_6	52
$Fe(acac)_3$	**C_6H_6**	**74**
$Fe(acac)_3$	THF	36

Making Organosodium Chemistry Even More Sustainable






Nat. Synth. **2025**, 10.1038/s44160-025-00771-1

Fe-catalysed Cross-coupling of Organosodium Reagents


UNIVERSITÄT BERN

Assessing cross-coupling reactivity



Catalyst	Yield A (%)	Yield B (%)
$Fe(acac)_3$	**83**	**12**
none	0	3
$Pd(acac)_2$	0	10
$Ni(acac)_2$	0	30
$Cu(acac)_2$	0	4


additive	Yield (%) (A:B ratio)
none	9 (30:70)
THF	29 (45:55)
15-crown-5	6 (11:89)
TMEDA	54 (76:24)
TMEDA (10 eq)	**96 (87:13)**

TMEDA = 


Fe is key to promote the C-C bond forming process

Nat. Synth. **2025**, 10.1038/s44160-025-00771-1

Fe-catalysed Cross-coupling of Organosodium Reagents



25 examples
Mild reaction conditions
No external ligands




Nat. Synth. **2025**, 10.1038/s44160-025-00771-1


Fe-catalysed Cross-coupling of Organosodium Reagents

UNIVERSITÄT BERN

Scope using primary alkyl bromides



Sterospecific coupling with cholesteryl bromide




Nat. Synth. **2025**, 10.1038/s44160-025-00771-1


Fe-catalysed Cross-coupling of Organosodium Reagents

UNIVERSITÄT BERN

Understanding the role of TMEDA




deaggreagation and activation of NaAr




Coordinative stabilisation of Fe(II) center



Andreu Tortajada







Nat. Synth. **2025**, 10.1038/s44160-025-00771-1

Fe-catalysed Cross-coupling of Organosodium Reagents

Radical-Clock Experiments:




Consistent with alkyl radical formation




M. Neidig, *Angew. Chem. Int. Ed.* **2022**, *61*, e202114986:
Fe cataysed cross-coupling of ArMgX

Nat. Synth. **2025**, 10.1038/s44160-025-00771-1


Harnessing Organopotassium Reagents for Cross-coupling Applications




Abundance: 0.002 %
Electronegativity: 0.98
Size: 145 pm



Abundance: 2.6 %
Electronegativity: 0.93
Size: 180 pm



Abundance: 2.4 %
Electronegativity: 0.82
Size: 235 pm




$[KCH_2SiMe_3(PMDETA)]_n$

Organopotassium Reagents

- highly aggregated structures
- limited solubility in hydrocarbons
- untamed reactivity
- no applications in cross-coupling reactions

R. E. Mulvey, *Eur. J. Inorg. Chem.* **2011**, 721

Assessing ability of organopotassium reagents for C-H metalation




KR is drastically superior as a base!

MR	Yield (%)
KCH_2SiMe_3	**80**
*n*BuLi + K*O*t*Bu*	65
$LiCH_2SiMe_3$	0
$NaCH_2SiMe_3$	5




KCH_2Ph generated *in situ in hexane*

Harnessing Organopotassium Reagents for Cross-coupling Applications



1 eq. 1 eq.


Pd cat	Yield (%)
none	5
Pd(P*t*Bu$_3$)$_2$	4



one pot approach:
$LiCH_2SiMe_3$ / KO*t*Bu
multigram synthesis:
85% yield (7 mmol scale)


Pd(P*t*Bu$_3$)$_2$

cross couplings of **Li**Ar with **ArBr**:
Feringa et al, *Nat. Chem.* **2013**, *5*, 667



Pd-PEPPSI-IPent

cross couplings of **Na**Ar with **ArBr**:
Asako et al, *Nat. Cat.* **2019**, *2*, 297



P1
pinkYPhos-Pd(Ind)Cl

cross couplings of **Li**Ar with **ArCl**:
Gessner et al, *Angew. Chem. Int. Ed.* **2020**, *59*, 20596

highly electron rich ylide phosphine



David Anderson




Daniel Knyszek





Viktoria Gessner


J. Am. Chem. Soc., **2025**, 147, 5417


Harnessing Organopotassium Reagents for Cross-coupling Applications


































- direct metalation
- fast catalysis
- over 25 examples
- simple one-pot, RT approach





























J. Am. Chem. Soc., **2025**, 147, 5417


Harnessing Organopotassium Reagents for Cross-coupling Applications





further potassium reagents




















From 

pK_A = 43







Limitations




7 % unselective cross-coupling




12 %, unselective cross-coupling




inaccessible



inaccessible




inaccessible





10 % unselective metallation

J. Am. Chem. Soc., **2025**, 147, 5417

Conclusions and Outlook









- ✓ power of Na and K to mediate metalation processes
- ✓ application in catalysis
- ✓ reactivity underpinned by **coordination effects** and **chemical cooperation**!

Angew. Chem. Int. Ed.,**2024**, 63, e202313556 (*Minireview*)

*n*BuNa and NaTMP: synthesis in a gram scale




Commercial reagents ▪ *Multi gram scale* ▪ *High purity*








Andreu Tortajada,
David Anderson

UNIVERSITÄT BERN









Helv. Chim. Acta **2022**, *105*, e202200060 (*open access*)

Acknowledgements

EH group

Jennifer Kuziola
Luca Vedani
Na Jin
Jelena Krizic
Rafela Feuz
Zippora Brüllhardt
Ioannis Vagiakos
Talia Mathers
Simon Kanbrock

David Sanchez
Roel Bienenmann
Sophia Belrhomari
Leon Liu
Clarence Tan
Andy Arribas-Domingo

Former members:
Andreu Tortajada
David Anderson

Collaborators

Robert Mulvey (Strathclyde, UK)
Laura Torrente (Cambridge, UK)
Karen Edler (Lund, Sweden)
Simon Grabowsky (Bern, Switzerland)
Sobi Asako (RIKEN, Japan)
Laurean Ilies (RIKEN, Japan)
Joaquin Garcia-Alvarez (Oviedo Spain)
Vito Capriati (Bari, Italy)
Dietmar Stalke (Göttigen, Germany)
Viktoria Gessner (Ruhr University)
Marie-Eve Perrin (Lyon, France)
Max Garcia-Melchor (Spain)
Karsten Meyer (Erlangen, Germany)
Odile Eisenstein (Montpellier, France)
Michele Cascella (Oslo, Norway)
Jesus Campos (IIQ, Spain)
David Balcells (Oslo, Norway)



Hevia Group

FNSNF

Alexander von HUMBOLDT STIFTUNG

Swiss Government Excellence Scholarship

frA FUNDACIÓN RAMÓN ARECES

@EvaHeviaGroup.bsky.social

Walter Benjamin-Programm DFG Deutsche Forschungsgemeinschaft